

## References

1. British Orthopaedic Association (BOA). The Management of Patients with Pelvic Fractures. British Orthopaedic Association Standards for Trauma and Orthopaedics (BOASTs). British Orthopaedic Association; 2018. Accessed December 1, 2023. <https://www.boa.ac.uk/standards-guidance/boasts.html>
2. Cullinane DC, Schiller HJ, Zielinski MD, et al. Eastern Association for the Surgery of Trauma practice management guidelines for hemorrhage in pelvic fracture--update and systematic review. *J Trauma*. Dec 2011;71(6):1850-68. doi:10.1097/TA.0b013e31823dca9a
3. Coccolini F, Stahel PF, Montori G, et al. Pelvic trauma: WSES classification and guidelines. *World J Emerg Surg*. 2017;12:5. doi:10.1186/s13017-017-0117-6
4. Yeung LL, McDonald AA, Como JJ, et al. Management of blunt force bladder injuries: A practice management guideline from the Eastern Association for the Surgery of Trauma. *J Trauma Acute Care Surg*. Feb 2019;86(2):326-336. doi:10.1097/ta.0000000000002132
5. British Orthopaedic Association (BOA). The Management of Urological Trauma Associated with Pelvic Fractures. British Orthopaedic Association Standards for Trauma and Orthopaedics (BOASTs). British Orthopaedic Association; 2016. Accessed December 1, 2023. <https://www.boa.ac.uk/standards-guidance/boasts.html>
6. Morey AF, Brandes S, Dugi DD, 3rd, et al. Urotrauma: AUA guideline. *J Urol*. Aug 2014;192(2):327-35. doi:10.1016/j.juro.2014.05.004
7. Black SR, Sathy AK, Jo C, Wiley MR, Minei JP, Starr AJ. Improved Survival After Pelvic Fracture: 13-Year Experience at a Single Trauma Center Using a Multidisciplinary Institutional Protocol. *J Orthop Trauma*. Jan 2016;30(1):22-8. doi:10.1097/bot.0000000000000443
8. Biffi WL, Smith WR, Moore EE, et al. Evolution of a multidisciplinary clinical pathway for the management of unstable patients with pelvic fractures. *Ann Surg*. Jun 2001;233(6):843-50. doi:10.1097/00000658-200106000-00015
9. Cheng M, Cheung MT, Lee KY, et al. Improvement in institutional protocols leads to decreased mortality in patients with haemodynamically unstable pelvic fractures. *Emerg Med J*. Mar 2015;32(3):214-20. doi:10.1136/emermed-2012-202009
10. Balogh Z, Caldwell E, Heetveld M, et al. Institutional practice guidelines on management of pelvic fracture-related hemodynamic instability: do they make a difference? *J Trauma*. Apr 2005;58(4):778-82. doi:10.1097/01.ta.0000158251.40760.b2
11. Vernamonti JP, Holcomb J, Mick NW, et al. 'Step Up' approach to the application of REBOA technology in a rural trauma system. *Trauma Surg Acute Care Open*. 2019;4(1):e000335. doi:10.1136/tsaco-2019-000335
12. DuBose JJ, Scalea TM, Brenner M, et al. The AAST prospective Aortic Occlusion for Resuscitation in Trauma and Acute Care Surgery (AORTA) registry: Data on contemporary utilization and outcomes of aortic occlusion and resuscitative balloon occlusion of the aorta (REBOA). *J Trauma Acute Care Surg*. Sep 2016;81(3):409-19. doi:10.1097/ta.0000000000001079

13. Gamberini E, Coccolini F, Tamagnini B, et al. Resuscitative Endovascular Balloon Occlusion of the Aorta in trauma: a systematic review of the literature. *World J Emerg Surg.* 2017;12:42. doi:10.1186/s13017-017-0153-2
14. Gardner MJ, Chip Routt ML, Jr. The antishock iliosacral screw. *J Orthop Trauma.* Oct 2010;24(10):e86-9. doi:10.1097/BOT.0b013e3181c81d65
15. Miller PR, Moore PS, Mansell E, Meredith JW, Chang MC. External fixation or arteriogram in bleeding pelvic fracture: initial therapy guided by markers of arterial hemorrhage. *J Trauma.* Mar 2003;54(3):437-43. doi:10.1097/01.Ta.0000053397.33827.Dd
16. Hymel A, Asturias S, Zhao F, et al. Selective versus nonselective embolization versus no embolization in pelvic trauma: A multicenter retrospective cohort study. *J Trauma Acute Care Surg.* Sep 2017;83(3):361-367. doi:10.1097/ta.0000000000001554